



## AMENDMENTS TO THE CLAIMS


1. (original) A method for the synthesis of severely sterically hindered secondary aminoether alcohols of the formula



wherein  $R^1$  and  $R^2$  are each selected from the group consisting of alkyl, hydroxyalkyl radicals having 1 to 4 carbon atoms or in combination with the carbon atom to which they are attached they form a cycloalkyl group having 3 to 8 carbon atoms, and  $R^3$  is selected from the group consisting of hydrogen, alkyl hydroxyalkyl radicals having 1 to 4 carbon atoms, and mixtures thereof, and  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$  and  $R^{11}$  are the same or different and are selected from the group consisting of hydrogen, alkyl and hydroxyalkyl radicals having 1 to 4 carbons provided that at least one of  $R^4$  or  $R^5$  bonded to the carbon atom directly bonded to the nitrogen atom is an alkyl or hydroxyalkyl radical when  $R^3$  is hydrogen, the process involving reacting an organic carboxylic acid or salt of a carboxylic acid of the formula




wherein  $R^{12}$  is selected from the group consisting of alkyl radicals having 1 to 4 carbon atoms, aryl radicals bearing hydrogen or one or more  $C_1$ - $C_{10}$  alkyl groups substituted thereon, and mixtures thereof, and Y is selected from the group consisting of hydrogen, alkali metal, ammonium, and mixtures thereof, with a sulfonyl halide, a sulfonyl halide, a mixed sulfonyl ester halide, or a mixed sulfonyl amide halide of the formula




wherein X is selected from the group consisting of F, Cl, Br, I, and mixtures thereof, and  $R^{14}$  and  $R^{14'}$  are the same or different and each is selected from the group consisting of alkyl radicals having 1 to 4 carbon atoms, haloalkyl radicals of the formula  $C_nH_{(2n+1)-w}Z_w$


wherein n is 1 to 4, Z is selected from the group consisting of F, Cl, Br, I, and mixtures thereof, and w ranges from 1 to 5, and aryl radicals




wherein R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup>, and R<sup>19</sup> are the same or different and are selected from hydrogen and alkyl radicals having 1 to 20 carbon atoms, and mixtures thereof, to yield acyl sulfonate material of the general formula





which is then reacted with a dioxane of the formula




wherein R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, and R<sup>11</sup> are the same or different and are selected from hydrogen, alkyl and hydroxyalkyl radicals having 1 to 4 carbons to yield







or




which is then aminated with an alkylamine of the formula



wherein R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> are as previously defined to yield



which is then hydrolyzed with base to yield



2. (original) The method of claim 1 for the synthesis of severely sterically hindered secondary aminoether alcohols using sulfonyl halide of the formula R<sup>14</sup>SO<sub>2</sub>X.

3. (original) The method of claim 1 for the synthesis of severely sterically hindered secondary aminoether alcohols using sulfuryl halide of the formula  $\text{SO}_2\text{X}_2$ .

4. (original) The method of claim 1 for the synthesis of severely sterically hindered secondary aminoether alcohols using the mixed sulfuryl ester halide of the formula  $\text{R}^{14}\text{OSO}_2\text{X}$ .

5. (original) The method of claim 1 for the synthesis of severely sterically hindered secondary aminoether alcohols using the mixed sulfuryl amide halide of the formula  $\text{R}_2^{14,14'}\text{NSO}_2\text{X}$ .

6. (currently amended) The method of according to ~~any one of the preceding claims~~ claim 1, 2, 3, 4 or 5 wherein  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  are methyl radicals.

7. (currently amended) The method of according to ~~any one of the preceding claims~~ claim 1, 2, 3, 4 or 5 wherein  $\text{R}^4$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$ ,  $\text{R}^8$ ,  $\text{R}^9$ ,  $\text{R}^{10}$ , and  $\text{R}^{11}$  are hydrogen.

8. (currently amended) The method of according to ~~any one of the preceding claims~~ claim 1, 2, 3, 4 or 5 wherein  $\text{R}^{15}$ ,  $\text{R}^{16}$ ,  $\text{R}^{18}$ , and  $\text{R}^{19}$  are hydrogen and  $\text{R}^{17}$  is hydrogen or methyl.

9. (currently amended) The method of according to ~~any one of the preceding claims~~ claim 1, 2, 3, 4 or 5 wherein the base is selected from alkali metal hydroxide, alkali metal alkoxide, or alkali metal carbonate.

10. (currently amended) The method of according to ~~any one of the preceding claims~~ claim 1, 2, 3, 4 or 5 wherein Y is hydrogen or sodium.

11. (currently amended) The method of according to ~~any one of the preceding claims~~ claim 1, 2, 3, 4 or 5 wherein  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  are methyl,  $\text{R}^4$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$ ,  $\text{R}^8$ ,  $\text{R}^9$ ,  $\text{R}^{10}$ , and  $\text{R}^{11}$  are hydrogen,  $\text{R}^{15}$ ,  $\text{R}^{16}$ ,  $\text{R}^{18}$ , and  $\text{R}^{19}$  are hydrogen,  $\text{R}^{17}$  is hydrogen or methyl, and Y is hydrogen, sodium, or ammonium.

12. (currently amended) The method of according to ~~any one of the preceding claims~~ claim 1, 2, 3, 4 or 5 wherein the acyl sulfonate is made by reacting organic carboxylic acid or the salt of a carboxylic acid with the sulfonyl halide, sulfuryl halide, mixed sulfuryl ester halide or mixed sulfuryl amide halide at a temperature in the range of between about -20 to 200°C at a pressure between about 1 bar and 100 bars, the acyl sulfonate is reacted with the dioxane at a molar ratio of dioxane to acyl sulfonate in the range of 1:1 to 10:1 at a temperature of between about 50°C to about 200°C to yield a cleavage product, the cleavage product and the alkyl amine reacted at an amine to sulfonate group ratio ranging from about stoichiometric to about 10:1 at pressure of from about atmospheric (1 bar) to about 100 bars at temperature of from about 40°C to about 200°C, and the resulting aminated product is hydrolyzed with base at a temperature from about 20°C to about 110°C.

13. (currently amended) The method of according to ~~any one of the preceding claims~~ claim 1, 2, 3, 4 or 5 wherein the organic carboxylic acid or the salt thereof, the sulfonyl halide, sulfuryl halide, mixed sulfuryl ester halide or mixed sulfuryl amide halide and the dioxane are combined in a single step to produce a reaction mixture, the reaction mixture being heated at a temperature of between about 50°C to about 200°C to produce the cleavage product, the cleavage product and the alkylamine are reacted at an amine to cleavage product ratio ranging from about stoichiometric to about 10:1 at a pressure from about atmospheric (1 bar) to about 100 bars at a temperature of from about 40°C to about 200°C, the resulting aminated product being reacted with base at a temperature from about 20°C to about 110°C.

U.S. National Phase of PCT/US2005/003052  
Preliminary Amendment  
Family Number: P2004J004

Applicants request that the present application be examined on the basis of the above presented amended claims.

Respectfully submitted,



Joseph J. Allocca

Attorney for Applicants

Registration No. 27,766

Telephone Number: (908) 730-3629

Facsimile Number: (908) 730-3649

☒ Pursuant to 37 CFR 1.34(a)

ExxonMobil Research and Engineering Company  
P. O. Box 900  
Annandale, New Jersey 08801-0900

JJA:kak  
7/18/06